

COMPACTNESS OF THE SPACE OF GENUS-ONE HELICOIDS

JACOB BERNSTEIN AND CHRISTINE BREINER

ABSTRACT. Using the lamination theory developed by Colding and Minicozzi for sequences of embedded, finite genus minimal surfaces with boundaries going to infinity [5], we show that the space of genus-one helicoids is compact (modulo rigid motions and homotheties). This generalizes a result of Hoffman and White [12].

1. INTRODUCTION

We investigate the finer geometric structure of complete, one-ended, genus- g minimal surfaces embedded in $\mathbb{R}^3$. Recall that Corollary 1.3 in [2] determined that every such surface, the space of which we denoted by $\mathcal{E}(1, g)$, is asymptotic to a helicoid and hence the terminology “genus- g helicoid” is warranted. We approach this by showing certain compactness properties for $\mathcal{E}(1, g)$ which will allow us to further bound the geometry of genus- g helicoids. For technical reasons, it is much simpler to treat the case $g = 1$, which we do in this paper. The compactness properties of $\mathcal{E}(1, g)$ for $g \geq 2$ will be discussed in [1].

Our main result shows that (after suitably normalizing) the space of genus-one helicoids is compact. This generalizes a result of Hoffman and White [12] in which they show compactness for the space of symmetric genus-one helicoids (i.e. genus-one helicoids that contain two coordinate axes). Our result establishes a connection between the asymptotic geometry of a genus-one helicoid and the geometry of the handle. Whether such compactness holds in $\mathcal{E}(1, g)$ when $g \geq 2$ is still unclear.

Theorem 1.1. *Let $\Sigma_i \in \mathcal{E}(1, 1)$ with each Σ_i asymptotic to H , a fixed helicoid. Then, a sub-sequence converges uniformly in C^∞ on compact subsets of $\mathbb{R}^3$ with multiplicity one to $\Sigma_\infty \in \mathcal{E}(1, 1) \cup \{H\}$ with Σ_∞ asymptotic to (or equaling) H .*

Remark 1.2. Translations along the axis show how H may occur as a limit.

To prove this, we develop a more general compactness result used also in [1]. We consider a sequence $\Sigma_i \in \mathcal{E}(1, g, R_i)$, where $R_i \rightarrow \infty$, that has uniform extrinsic control on the position and size of the genus. Here $\mathcal{E}(e, g, R)$ denotes the set of smooth, connected, properly embedded minimal surfaces, $\Sigma \subset \mathbb{R}^3$, so that Σ has genus g and $\partial\Sigma \subset \partial B_R(0)$ is smooth and has e components. We show the sequence Σ_i has uniformly bounded curvature and so contain a convergent sub-sequence.

Theorem 1.3. *Suppose $\Sigma_i \in \mathcal{E}(1, g, R_i)$ are such that $1 = r_-(\Sigma_i) \geq \alpha r_+(\Sigma_i)$, the genus of each Σ_i is centered at 0 and $R_i \rightarrow \infty$. Then a sub-sequence of the Σ_i converges uniformly in C^∞ on compact subsets of $\mathbb{R}^3$ and with multiplicity one to a surface $\Sigma_\infty \in \mathcal{E}(1, g)$ and $1 = r_-(\Sigma_\infty) \geq \alpha r_+(\Sigma_\infty)$.*

The first author was supported in part by NSF grant DMS 0606629.

For the meaning of $r_+(\Sigma), r_-(\Sigma)$, see Definitions 3.1 and 3.3. Roughly, $r_+(\Sigma)$ measures the extrinsic spread of the handles of Σ , while $r_-(\Sigma)$ measures the size of the smallest handle. When $g = 1$, as $r_-(\Sigma) = r_+(\Sigma)$, rescaling arguments make Theorem 1.3 particularly strong. For example, it allows one to give a description of elements of $\mathcal{E}(1, 1, R)$ which have $r(\Sigma)$ small relative to R . This is Theorem 4.3, which says such a surface is close, in a Lipschitz sense, to a piece of a genus-one helicoid. This is comparable to the description of the shape of an embedded minimal disks near a point of large curvature given by Theorem 1.1 in [3].

The proof of Theorem 1.3 uses extensively the structural theory for embedded minimal surfaces of Colding and Minicozzi [5, 6, 7, 8, 9]. In particular, we make use of three important consequences of their work: the one-sided curvature estimates – Theorem 0.2 of [9]; the chord-arc bounds for minimal disks – Proposition 1.1 of [10]; and, most importantly, their lamination theory for finite genus surfaces – Theorem 0.9 of [5]. As our proof depends most critically on this last result, we discuss it in some detail in Appendix A below. In particular, we summarize Theorem 0.9 of [5] as Theorem A.3.

Let us briefly outline the structure of the paper. In Section 2, we prove uniform curvature bounds for sequences of minimal surfaces from a larger class than that of Theorem 1.3. These surfaces have uniformly finite topology normalized so that, intrinsically, the topology does not concentrate or disappear. As a consequence, we deduce smooth sub-sequential convergence to a complete, embedded minimal surface in $\mathbb{R}^3$. This result is more general than Theorem 1.3, and consequently there are more possibilities for the limit surfaces. While it is implicit in work of Meeks, Perez and Ros [13, 14], we give our own proof:

Theorem 1.4. *Let $\Sigma_i \in \mathcal{E}(e, g, R_i)$ ($e, g \geq 1$) be such that $0 \in \Sigma_i$, $\text{inj}_{\Sigma_i} \geq 1$, $\text{inj}_{\Sigma_i}(0) \leq \Delta$ and $R_i \rightarrow \infty$. Then a sub-sequence of the Σ_i converge smoothly on compact subsets of $\mathbb{R}^3$ with multiplicity one to a non-simply connected minimal surface in $\cup_{1 \leq k \leq e+g, 0 \leq l \leq g} \mathcal{E}(k, l)$.*

We will prove Theorem 1.4 by using the lamination theory of Colding and Minicozzi. The key fact is that the weak chord-arc bounds (i.e. Proposition 1.1 of [10]) allow us to show that the sequence Σ_i is ULSC, Uniformly Locally Simply Connected (see Definition A.1 below). That is, there is a small, but uniform, *extrinsic* scale on which the sequence is simply connected (the uniform lower bounds for the injectivity radius provide such a uniform *intrinsic* scale). This allows the local application of the work of Colding and Minicozzi for disks.

The uniform upper bound on the injectivity radius at 0 implies the existence of a closed geodesic, γ_i , in each Σ_i , close to 0 and with uniformly bounded length. Using the γ_i and the lamination theorem, we establish uniform curvature bounds on compact subsets of $\mathbb{R}^3$ for the sequence. Indeed, suppose one had a sequence with curvature blowing-up; then a sub-sequence would converge to a singular lamination as in [5]. The nature of the convergence implies that any sequence of closed geodesics in the surfaces, which lie in a fixed extrinsic ball, must converge (in a Hausdorff sense) to a subset of the singular axis; in particular this is true of the γ_i . However, this must contradict certain chord-arc properties of the surfaces and thus cannot occur. The uniform curvature bounds, together with Schauder estimates and the Arzela-Ascoli theorem then prove Theorem 1.4.

In Section 3.2, we use Theorem 1.4 to deduce that, for sequences in $\mathcal{E}(1, g, R_i)$, as long as the genus stays inside a fixed extrinsic ball and does not shrink off, then

one has convergence to an element of $\mathcal{E}(1, g)$. Indeed, with such uniform control, the no-mixing theorem of [5] gives a uniform lower bound on the injectivity radius and so Theorem 1.4 applies; that the genus remains in a fixed ball implies that the limit surface must belong to $\mathcal{E}(1, g)$. The proof of this gives Theorem 1.3. In order to show our main application, i.e. Theorem 1.1, which we prove in Section 4.1, we couple Theorem 1.3 with the fact that the surfaces are asymptotic to helicoids. The connection between the convergence on compact subsets of $\mathbb{R}^3$ and the asymptotic behavior at the end is made using the holomorphic Weierstrass data.

Throughout, we denote balls in $\mathbb{R}^3$, centered at x with radius r , by $B_r(x)$, while intrinsic balls in a surface, Σ , are denoted by $\mathcal{B}_r^\Sigma(x)$. The norm squared of the second fundamental form of Σ is $|A|^2$. At various points we will need to consider $\Sigma \cap B_r(x)$ and when we do so we always assume $\partial B_r(x)$ meets Σ transversely.

2. CURVATURE BOUNDS AT THE SCALE OF THE TOPOLOGY

As mentioned in the introduction, the lamination theory of [5] proves crucial for the proof of Theorem 1.4. Indeed, this implies that there is a sub-sequence with either uniform curvature bounds or one of two possible singular convergence models. In the latter case, a simple topological argument will rule out one singular model and so imply that the sequence behaves like the blow-down of a helicoid – i.e. like Theorem 0.1 of [9]. This contradicts the upper bound on the injectivity radius at the origin, which proves the desired curvature bounds.

2.1. Technical lemmas. In order to obtain the curvature bounds, we will need four technical lemmas. We first note the following simple topological consequence of the maximum principle:

Proposition 2.1. *Let $\Sigma \in \mathcal{E}(e, g, R)$ and suppose $B_r(x) \subset B_R$. Then, for any component Σ_0 of $\Sigma \cap B_r(x)$, $\partial \Sigma_0$ has at most $g + e$ components.*

Proof. Let Σ_i , $1 \leq i \leq n$, be the components of $\Sigma \setminus \Sigma_0$. The Σ_i are smooth compact surfaces with boundary, and so $\partial \Sigma_i$ is a finite collection of circles. Thus, the Euler characteristic satisfies $\chi(\Sigma) = \sum_{i=0}^n \chi(\Sigma_i)$. By the classification of surfaces, $\chi(\Sigma) = 2 - 2g - e$ and $\chi(\Sigma_i) = 2 - 2g_i - e_i$ where g_i is the genus of Σ_i and e_i number of components of $\partial \Sigma_i$. Note that, $\sum_{i=0}^n g_i \leq g$ and $\sum_{i=1}^n e_i = e_0 + e$. Thus, we compute that $e_0 = n + g - \sum_{i=0}^n g_i$. The maximum principle implies that $n \leq e$ (as any Σ_i , for $i \geq 1$, must meet ∂B_R). Thus, $e_0 \leq e + g$. $\square$

Next we note it is impossible to minimally immerse a (intrinsically) long and thin cylinder in $\mathbb{R}^3$:

Lemma 2.2. *Let Γ be a minimal surface with genus g and with $\partial \Gamma = \gamma_1 \cup \gamma_2$ where the γ_i are smooth and satisfy $\int_{\gamma_i} 1 + |k_g| \leq C_1$. Then, there exists $C_2 = C_2(g, C_1)$ so that $\text{dist}_\Gamma(\gamma_1, \gamma_2) \leq C_2$.*

Proof. It is clear that there are points $p_1, p_2 \in \mathbb{R}^3$ so that $\gamma_i \subset B_{C_1}(p_i)$. Thus, up to rotating and translating Γ , $\gamma_i \subset \{x_1^2 + x_2^2 \leq C_1^2\}$ and so by the convex hull property, Γ itself lies in this cylinder. Notice that $|\nabla \text{dist}_{\mathbb{S}^2}(\mathbf{n}, \mathbf{n}_0)| \leq |A|$ (here $\mathbf{n}$ is the normal to Γ) and so there exists a uniform constant δ_0 so that if $s \sup_{\mathcal{B}_s(p)} |A| \leq \delta_0$ then $\mathcal{B}_s(p)$ can be expressed as a graph over $T_p \Gamma$ where the graph has gradient $\leq 1/100$. Indeed, for such a p if $q \in \mathcal{B}_s(p)$ then $|q - p| \geq \frac{9}{10} \text{dist}_\Gamma(p, q)$. We refer the reader to Section 2 of [7] for more details. Thus, if $p \in \Gamma$ with $\mathcal{B}_{4C_1}(p) \cap \partial \Gamma = \emptyset$ and

$4C_1 \sup_{\mathcal{B}_{4C_1}(p)} |A| \leq \delta_0$ then $\mathcal{B}_{4C_1}(p)$ cannot lie in the cylinder $\{x_1^2 + x_2^2 \leq C_1^2\}$. We claim that if $\text{dist}_\Gamma(\gamma_1, \gamma_2)$ was very large we could find such a p yielding the desired contradiction.

We use the intrinsic version of a result of Choi-Schoen [4] on Γ . That is, there exists a universal $\epsilon > 0$ so that if $\mathcal{B}_s(p) \subset \Gamma$ is disjoint from $\partial\Gamma$ and $\int_{\mathcal{B}_s} |A|^2 \leq \delta\epsilon$ then $\sup_{\mathcal{B}_{s-t}(p)} t^2 |A|^2 \leq \delta$. We emphasize that s is not a priori restricted to be smaller than $\text{inj}_\Gamma(p)$. By the Gauss-Bonnet theorem, $\int_\Gamma |A|^2 \leq 8\pi g + 4C_1 = C'_1$. If $\text{dist}_\Gamma(\gamma_1, \gamma_2) \geq \frac{32C'_1 C_1}{\delta_0^2 \epsilon}$ then by the pigeonhole principle, there is a $p \in \Gamma$ so that $\mathcal{B}_{8C_1}(p) \cap \partial\Gamma = \emptyset$ and $\int_{\mathcal{B}_{8C_1}(p)} |A|^2 \leq \delta_0^2 \epsilon$. The Choi-Schoen theorem and the conclusion of the preceding paragraph then imply that $C_2 \leq \frac{32C'_1 C_1}{\delta_0^2 \epsilon}$. $\square$

We need that minimizing geodesics in almost flat surfaces are almost straight:

Lemma 2.3. *Let $u : D_{3/2}(0) \rightarrow \mathbb{R}$ and let Σ be the graph of u . Then for any $\delta > 0$ there is an $\epsilon > 0$ so: if $\|u\|_{C^2} \leq \epsilon$ and $\gamma \subset \Sigma$ a minimizing geodesic with $\partial\gamma = \{p_-, p_+\} \in \partial B_1(0)$ satisfies $\gamma \cap B_\epsilon(0) \neq \emptyset$, then there is a line $0 \in L$ so that the Hausdorff distance between γ and $L \cap D_1(0)$ is less than δ .*

Proof. Fix $\delta > 0$ and suppose this result was false. Then there is a sequence of Σ_i and u_i with $u_i \rightarrow 0$ in C^2 and points $p_\pm^i$ connected by minimizing geodesic $\gamma_i \subset \Sigma_i$ meeting $B_{\epsilon_i}(0)$ where $1 > \epsilon_i \rightarrow 0$, but the conclusion of the lemma does not hold.

Chord-arc bounds for graphs with small C^1 bounds imply $\text{dist}_{\mathbb{R}^3}(p_+^i, p_-^i) \geq 1/2$. Letting $u_i \rightarrow 0$, we get $p_\pm^i \rightarrow p_\pm^\infty \in \partial D_1(0)$ and $|p_-^\infty - p_+^\infty| \geq 1/2$. Now, let L be the line connecting $p_\pm^\infty$. We claim that $0 \in L$. If not, then $\ell(L \cap D_1(0)) = 2 - 4\alpha$ for some $\alpha > 0$. Let L_i be the graph (of u_i) over $L \cap D_1(0)$, so L_i is a segment that is a subset of Σ_i . Clearly, for some i_0 , $i \geq i_0$ implies $\text{dist}_{\Sigma_i}(p_\pm^i, L_i) < \alpha$ and $\ell(L_i) < \ell(L \cap D_1(0)) + \alpha$; and thus, $\ell(\gamma_i) < 2 - \alpha$. On the other hand, the hypotheses give a $p^i \in \gamma_i$ with $p^i \rightarrow 0$. Thus, by possibly increasing i_0 , for $i \geq i_0$, $\ell(\gamma_i) \geq 2 - \alpha$. Similar arguments show, for i large, $L \cap D_1(0)$ must be Hausdorff close to γ_i , yielding the desired contradiction. $\square$

Our final technical lemma shows that, for a ULSC sequence Σ_i converging to a minimal lamination with a single singular line, any closed geodesics in the Σ_i that lie in a fixed extrinsic ball must collapse to the singular set.

Lemma 2.4. *Fix $\Delta > 0$. Suppose $\Sigma_i \in \mathcal{E}(e, g, R_i)$, $R_i \rightarrow \infty$, the Σ_i converge to the singular lamination $\mathcal{L}$ with singular set $\mathcal{S} = \mathcal{S}_{\text{ulsc}}$ in the sense of Theorem 0.9 of [5], and that $\mathcal{S}$ is the x_3 -axis. Then, given $\epsilon > 0$ there exists an i_0 so that for $i \geq i_0$, if γ_i is an embedded closed geodesic in Σ_i , with $\gamma_i \subset B_\Delta$, then $\gamma_i \subset T_\epsilon(\mathcal{S})$, the extrinsic ϵ -tubular neighborhood of $\mathcal{S}$.*

Proof. Suppose the lemma was not true; then there exists a sub-sequence of the Σ_i so that γ_i intersects $K_\epsilon = \overline{B}_\Delta(0) \setminus T_\epsilon(\mathcal{S})$. Choose $p_i \in \gamma_i \cap \overline{K}_\epsilon$ so that $x_1^2 + x_2^2 = \rho^2$ achieves its maximum on γ_i at p_i . After possible passing to a further sub-sequence, $p_i \rightarrow p_\infty \in \overline{K}_\epsilon$ with $\rho^2(p_\infty) \geq \epsilon$. Translate so that $x_3(p_\infty) = 0$. The convergence of [5] implies that for sufficiently large i , the component Γ_i of $B_{\epsilon/2}(p_\infty) \cap \Sigma_i$ containing p_i converges smoothly, as a graph, to the disk $B_{\epsilon/2}(p_\infty) \cap \{x_3 = 0\}$. Let γ'_i be the component of $B_{\epsilon/2}(p_\infty) \cap \gamma_i$ containing p_i with boundary points $q_i^\pm$.

For a given $\delta > 0$, there exists i large so that Γ_i , γ'_i satisfy the hypotheses of Lemma 2.3 (after a rescaling). Notice that for large i , Γ_i is very flat and in




FIGURE 1. The points of interest in the proof of Lemma 2.5.

particular is geodesically convex; hence γ'_i is the minimizing geodesic connecting $q_i^\pm$. If L_i is the line given by the lemma, γ'_i lies in the δ -tubular neighborhood of $L_i \cap D_{\epsilon/4}(p_\infty)$. By passing to a sub-sequence, the γ'_i converge to a segment of a line L in $\{x_3 = 0\}$ that goes through p_∞ . Because $\{x_3 = 0\}$ is transverse to the axis of the cylinder $\{y \in \mathbb{R}^3 | \rho^2(y) \leq \rho^2(p_\infty)\}$ and p_∞ is on the boundary of this cylinder, L cannot lie entirely within it. However, this implies there are points $q_i \in \gamma_i$ with $\lim_{i \rightarrow \infty} \rho^2(q_i) > \rho^2(p_\infty)$, giving a contradiction. $\square$

2.2. Proof of Theorem 1.4. We apply the preceding lemmas in order to show uniform curvature bounds. We note that similar techniques are used by Meeks and Rosenberg in [15] to achieve different results:

Lemma 2.5. *Let $\Sigma_i \in \mathcal{E}(e, g, R_i)$ be such that $0 \in \Sigma_i$, $\text{inj}_{\Sigma_i} \geq 1$, $\text{inj}_{\Sigma_i}(0) \leq \Delta$ and $R_i \rightarrow \infty$. Then a sub-sequence of the Σ_i satisfy, for any compact K ,*

$$(2.1) \quad \sup_i \sup_{K \cap \Sigma_i} |A|^2 < \infty.$$

Proof. If this was not the case then by the lamination theorem of [5] a sub-sequence of the Σ_i would converge to a singular lamination $\mathcal{L}$. For any $x \in \Sigma_i$ with $|x| \leq R_i/2$, the injectivity radius lower bound and the weak chord-arc bounds of [10] imply that there is a $\delta_0 > 0$ so the component of $B_{\delta_0}(x) \cap \Sigma_i$ containing x is a subset of $\mathcal{B}_{1/2}^{\Sigma_i}(x)$. For any fixed $x \in \mathbb{R}^3$, as long as i is large enough so $|x| \leq R_i/2$, this implies that every component of $B_{\delta_0}(x) \cap \Sigma_i$ is a disk. Thus, the sequence of Σ_i is ULSC and so $\mathcal{S} = \mathcal{S}_{ulsc}$. Hence, after rotating if needed, $\mathcal{L} = \{x_3 = t\}_{t \in \mathbb{R}}$ and $\mathcal{S}$ is parallel to the x_3 -axis and consists of either one or two lines. By Remark A.4 and Lemma 2.1 we see that $\mathcal{S}$ must consist of exactly one singular line. We next rule out any singular behavior. The injectivity bound at 0 and the non-positive curvature of Σ_i imply the existence of $0 \in \gamma'_i \subset \Sigma_i$, a geodesic lasso (i.e. a closed curve that is a geodesic but for at most one point) with $\ell(\gamma'_i) \leq 2\Delta$. The length bound implies $\gamma'_i \subset B_{3\Delta}$. The fact that Σ_i has non-positive curvature implies that γ'_i is not null-homotopic. We may minimize in the homotopy class of γ'_i to obtain a closed geodesic γ_i ; note that Lemma 2.2 allows us to do this even though Σ_i has boundary that is not a priori convex. Indeed, either γ'_i intersects γ_i and so $\gamma_i \subset B_{6\Delta}$ or, as $\ell(\gamma_i) \leq \ell(\gamma'_i)$,




FIGURE 2. Illustrating the consequence of the one sided curvature estimates

Lemma 2.2 implies that $\text{dist}_{\Sigma_i}(\gamma'_i, \gamma_i) \leq C$ for some large (but uniform) C , and so $\gamma_i \subset B_{(C+6)\Delta}(0)$.

Now for $\epsilon > 0$, let $T_\epsilon(\mathcal{S})$ be the extrinsic ϵ -tubular neighborhood of $\mathcal{S}$. Lemma 2.4 then implies there is an i_ϵ so that for all $i \geq i_\epsilon$, $\gamma_i \subset T_\epsilon(\mathcal{S})$.

For each i , fix $p_i \in \gamma_i$ so that $x_3(p_i) = \min \{x_3(p) : p \in \gamma_i\}$, i.e. the lowest point of γ_i . Then a sub-sequence of the p_i converge to $p_\infty \in \mathcal{S}$. Let p_+ be the point of intersection $\partial B_{\delta_0/2}(p_\infty) \cap \mathcal{S}$, chosen so $x_3(p_+) > x_3(p_\infty)$ with δ_0 as before. Pick i_0 large enough so for $i \geq i_0$, $|p_\infty - p_i| \leq \delta_0/4$. The choice of δ_0 implies that γ_i is not contained in $B_{\delta_0/2}(p_\infty)$; but for $i \geq i_0$, γ_i meets this ball. Let γ_i^0 be an arc of γ_i in $B_{\delta_0/2}(p_\infty)$ through p_i . Denote $\partial\gamma_i^0 = \{q_i^+, q_i^-\} \subset \partial B_{\delta_0/2}(p_\infty)$ (see Figure 1). For $i \geq i_0$, $\text{dist}_{\Sigma_i}(q_i^-, q_i^+) \geq \delta_0/3$. Indeed, for $i \geq i_0$, the length of γ_i^0 is bounded below by $\delta_0/2$, as $|q_i^\pm - p_i| \geq \delta_0/4$. On the other hand, the lower bound on the injectivity radius implies either $\text{dist}_\Sigma(q_i^-, q_i^+) > 1/2$ or both lie in a geodesically convex region with γ_i^0 the minimizing geodesic connecting them.

By Lemma 2.4, for any $\delta > 0$, there is an $i_\delta \geq i_0$ so for $i \geq i_\delta$, $q_i^\pm \in B_\delta(p_+)$. By the one-sided curvature estimate of [9], there exist $c > 1$ and $1 > \epsilon > 0$ so that if Σ_1, Σ_2 are disjoint embedded disks in B_{cR} with $\partial\Sigma_j \subset \partial B_{cR}$ and $B_{\epsilon R} \cap \Sigma_j \neq \emptyset$, then for all components Σ'_1 of $B_R \cap \Sigma_1$ that intersect $B_{\epsilon R}$, $\sup_{\Sigma'_1} |A|^2 \leq R^{-2}$. Thus, for δ such that $0 < \frac{2c}{\epsilon}\delta < \delta_0$, because $\lim_{i \rightarrow \infty} \sup_{\Sigma_i \cap B_\delta(p_+)} |A|^2 \rightarrow \infty$, there is an $i'_\delta \geq i_\delta$ so for $i \geq i'_\delta$ there is only one component of $B_{c\delta/\epsilon}(p_+) \cap \Sigma_i$ that meets $B_\delta(p_+)$. And thus, there is a $\sigma_i \subset B_{c\delta/\epsilon}(p_+) \cap \Sigma_i$ that connects $q_i^\pm$ (see Figure 2).

Choose $\gamma\delta = \frac{2c}{\epsilon}\delta < \delta_0$. Then, for $i \geq i'_\delta$, $q_i^- \in B_{\gamma\delta}(q_i^+)$, and thus the component Σ_i^δ of $B_{\gamma\delta}(q_i^+) \cap \Sigma_i$ that contains q_i^+ is a disk containing σ_i and thus q_i^- . Let $1 > \delta_1 > 0$ be given by the weak chord arc bounds (see [10]) and decrease δ , if necessary, so $\gamma\delta < \frac{1}{2}\delta_0\delta_1$. Then, $\mathcal{B}_{2\gamma\delta\delta_1^{-1}}^{\Sigma_i}(q_i^+)$ is a disk, and so $\Sigma_i^\delta \subset \mathcal{B}_{\gamma\delta\delta_1^{-1}}^{\Sigma_i}(q_i^+)$, by the weak chord-arc bounds. Thus, for $i > i'_\delta$ and δ sufficiently small, $\text{dist}_{\Sigma_i}(q_i^+, q_i^-) < \gamma\delta\delta_1^{-1} < \delta_0/3$, which gives a contradiction. $\square$

Theorem 1.4 will follow from these uniform curvature bounds as long as one further point is addressed. Namely, we must verify uniform area bounds for the sequence. For embedded minimal disks this follows immediately from Lemma 3.1 of [7]. For the more general surfaces we consider, one must use a type of chord-arc bound, the details of which are in Lemma C.3 of [5].

Proof. (Theorem 1.4) By Lemma 2.5, the curvature of the Σ_i is uniformly bounded on any compact subset of $\mathbb{R}^3$. Pick $R_j > \Delta$ with $R_j \rightarrow \infty$ and let Σ_i^j denote the connected component of $\Sigma_i \cap B_{R_j}(0)$ containing 0. Clearly, $\sup_{\Sigma_i^j} |A|^2 \leq C(j)$. For each j , Lemma C.3 of [5] implies the area of Σ_i^j is uniformly bounded. Hence, by standard compactness theory, for each j a sub-sequence of the Σ_i^j converge smoothly to an embedded, smooth minimal surface Σ_∞^j with $\partial \Sigma_\infty^j \subset \partial B_{R_j}$. Diagonalizing this sequence, gives a properly embedded surface Σ_∞ with $\Sigma_i^{j_i} \rightarrow \Sigma_\infty$.

Notice $\text{inj}_{\Sigma_i^j}(0) \leq \Delta$ implies $\text{inj}_{\Sigma_\infty}(0) \leq \Delta$, and thus Σ_∞ is not a disk. One may thus argue as in Appendix B of [5] to see that the $\Sigma_i^{j_i}$ converge to Σ_∞ with multiplicity 1. Roughly speaking, if the convergence was with a higher degree of multiplicity, one could construct a positive Jacobi function on Σ_∞ which would force Σ_∞ to be stable (by [11]) and hence be a plane by [16]. The convergence can only decrease the genus, so using Lemma 2.1 one sees $\Sigma_\infty \in \mathcal{E}(e', g')$ where $0 \leq g' \leq g$ and $0 \leq e' \leq e + g$. Finally, doing the same with other components of $B_{R_j} \cap \Sigma_i$ (if they exist) and noting that the only disjoint complete, properly embedded minimal surfaces of finite topology are parallel planes, one has $\Sigma_i \rightarrow \Sigma_\infty$. $\square$

3. PROOF OF THEOREM 1.3

3.1. Scale of the genus. We define the extrinsic scale(s) of the genus:

Definition 3.1. For $\Sigma \in \mathcal{E}(1, g, R)$ let

$$r_+(\Sigma) = \inf_{x \in B_R} \inf \{r : B_r(x) \subset B_R \text{ and } B_r(x) \cap \Sigma \text{ has a component of genus } g\}.$$

We call $r_+(\Sigma)$ the *outer extrinsic scale of the genus* of Σ . Furthermore, suppose for all $\epsilon > 0$, one of the components of $B_{r_+(\Sigma)+\epsilon}(x) \cap \Sigma$ has genus g ; then we say the genus is *centered at x* .

The outer scale of the genus measures how spread out all the handles are and the center of the genus should be thought of as a “center of mass” of the handles. We also need to measure the scale of individual handles and where the smallest handle (with respect to this scale) is located. To that end define:

Definition 3.2. For $\Sigma \in \mathcal{E}(1, g, R)$ and $x \in B_R(0)$ let

$$r_-(\Sigma, x) = \sup \{r : B_r(x) \subset B_R(0) \text{ and } B_r(x) \cap \Sigma \text{ is genus zero}\}.$$

If the genus of $B_r(x) \cap \Sigma$ is zero whenever $B_r(x) \subset B_R(0)$, set $r_-(\Sigma, x) = \infty$.

Definition 3.3. For $\Sigma \in \mathcal{E}(1, g, R)$ let

$$r_-(\Sigma) = \inf_{x \in B_R(0)} r_-(\Sigma, x).$$

We call $r_-(\Sigma)$ the *inner extrinsic scale of the genus* of Σ .

Remark 3.4. One easily checks that $R > r_+(\Sigma) \geq r_-(\Sigma)$ for $\Sigma \in \mathcal{E}(1, g, R)$, with equality holding if $g = 1$. When $g = 1$ we denote the common value by $r(\Sigma)$.

3.2. Two-sided bounds on the genus. Using the no-mixing theorem of [5], we show that uniform control on both the inner and outer extrinsic scales of the genus implies a uniform lower bound on the injectivity radius. By the weak chord-arc bounds of [10], uniform extrinsic control also gives a point with a uniform upper bound on its injectivity radius. We then apply Theorem 1.4 to obtain smooth sub-sequential convergence to some complete embedded minimal surface in $\mathbb{R}^3$. The uniform bound on the outer scale of the genus implies this limit must be a genus g -helicoid.

Lemma 3.5. *Fix $0 < \alpha \leq 1$ and $g \in \mathbb{Z}^+$. Then there exists $R_0 > 1$ and $1 > \delta_0 > 0$ depending only on α and g so: If $\Sigma \in \mathcal{E}(1, g, R)$ with $R \geq R_0$, $1 = r_-(\Sigma) \geq \alpha r_+(\Sigma)$, and the genus of Σ is centered at 0, then as long as $B_{\delta_0}(x) \subset B_R$, every component of $B_{\delta_0}(x) \cap \Sigma$ is a disk.*

Proof. Suppose the lemma was not true. Then, there exists a sequence of surfaces $\Sigma_i \in \mathcal{E}(1, g, R_i)$ with $1 = r_-(\Sigma_i) \geq \alpha r_+(\Sigma_i)$, $R_i \rightarrow \infty$, and the genus of Σ_i centered at 0. Further, there exist points x_i and a sequence $\delta_i \rightarrow 0$ so that one of the components of $B_{\delta_i}(x_i) \cap \Sigma_i$ is not a disk. Notice for fixed x and r , if $\bar{B}_r(x) \cap \bar{B}_{\alpha^{-1}}(0) = \emptyset$ with i large enough so $B_r(x) \subset B_{R_i}$, then each component of $B_r(x) \cap \Sigma_i$ is a disk. Thus, we may assume $x_i \in B_{3/2\alpha^{-1}}$. Because uniform curvature bounds would imply a uniform extrinsic scale on which Σ_i is graphical, we see that $\sup_{B_{2\alpha^{-1}} \cap \Sigma_i} |A|^2 \rightarrow \infty$. Hence, by Theorem 0.14 of [5], up to a sub-sequence, the Σ_i convergence to a singular lamination $\mathcal{L}$.

Let us now determine $\mathcal{L}$ and see that this gives a contradiction. By possibly passing to a further sub-sequence, we may assume that $x_i \rightarrow x_\infty$. Pick i_0 large enough so that $|x_i - x_\infty| + \delta_i \leq 1/8$ for all $i \geq i_0$. Then for $i \geq i_0$, $B_{\delta_i}(x_i) \subset B_{1/2}(x_\infty)$ and thus $B_{1/2}(x_\infty) \cap \Sigma_i$ contains a non-disk component. As $r_-(\Sigma_i) > 1/2$, this component has genus zero. Hence, the boundary of this component is not connected, and so x_∞ is a point of $\mathcal{S}_{neck}$ of the lamination $\mathcal{L}$ (see Definition A.2).

For any ball $B_r(x)$ with $\bar{B}_r(x) \cap \bar{B}_{\alpha^{-1}}(0) = \emptyset$ and i large enough so $B_r(x) \subset B_{R_i/2}(0)$, one has that all components of $B_r(x) \cap \Sigma_i$ are disks, and thus the maximum principle and the no-mixing theorem of [5] imply that the singular set $\mathcal{S}$ of $\mathcal{L}$ is contained in $B_{\alpha^{-1}}(0)$. As a consequence, one may rotate so that $\mathcal{L} \subset \{|x_3| \leq \alpha^{-1}\}$. Thus, for any $k \in \mathbb{N}$ there is an i_k so that for $i \geq i_k$, $\Sigma_i \cap B_k \subset \{|x_3| \leq 2\alpha^{-1}\}$ and $R_i > k^2$. Now set $\tilde{\Sigma}_k = \frac{1}{k}\Sigma_{i_k}$, so $\tilde{\Sigma}_k$ is a new sequence with the genus still centered at 0, $\partial\tilde{\Sigma}_k \cap B_k(0) = \emptyset$, $r_+(\tilde{\Sigma}_k) \leq \alpha^{-1}/k \rightarrow 0$, and $\tilde{\Sigma}_l \cap B_1 \subset \{|x_3| \leq 2(\alpha k)^{-1}\}$ for $l \geq k$. Clearly the curvature in $B_{2\alpha^{-1}}$ is still blowing up and so by possible passing to a sub-sequence we have convergence to a singular lamination $\tilde{\mathcal{L}}$. But $r_+(\tilde{\Sigma}_k) \rightarrow 0$ implies that $0 \in \mathcal{S}_{ulsc}$. However, $\tilde{\Sigma}_l \cap B_1$ converges (in a Hausdorff sense) to $\{x_3 = 0\} \cap B_1$, contradicting the convergence of Theorem 0.9 of [5]. $\square$

We now bound the injectivity radius of Σ above in terms of $r_+(\Sigma)$.

Lemma 3.6. *There exists R_0, Δ with $R_0 \geq 5\Delta > 10 > 0$, so: If $\Sigma \in \mathcal{E}(1, g, R)$, where $R > R_0$, and one of the components, Σ' , of $\Sigma \cap B_2(0)$ has positive genus, then the injectivity radius at all points of Σ' is bounded above by Δ .*

Proof. Pick $0 < \delta_1 \leq 1/2$ as in the weak chord-arc bounds of [10]. We claim that we may choose $\Delta = 4/\delta_1 \geq 3$ and $R_0 = 5\Delta$. To see this, suppose that Σ satisfies the hypotheses of the lemma for some $R > R_0$, and Σ' is a component of $\Sigma \cap B_2$ with positive genus, but the injectivity radius of some point $x \in \Sigma'$ is

(strictly) bounded below by Δ . Then $\mathcal{B}_{\Delta}^{\Sigma}(x)$ is disjoint from the boundary of Σ and is topologically a disk. Thus, the weak chord-arc bounds of [10] imply that the component of $B_4(x) \cap \Sigma$ containing x , $\Sigma_{x,4}$, is contained in the disk $\mathcal{B}_{\Delta/2}^{\Sigma}(x)$. The maximum principle implies $\Sigma_{x,4}$ is itself a disk; but $\Sigma' \subset \Sigma_{x,4}$, which gives a contradiction. $\square$

Corollary 3.7. *Fix $1 \geq \alpha > 0$ and let $\Sigma \in \mathcal{E}(1, g, R)$ and suppose that $1 = r_-(\Sigma) \geq \alpha r_+(\Sigma)$, $R \geq R_0 \alpha^{-1}$, the genus is centered at 0, and R_0, Δ are as above. Then there is a point $p_0 \in \Sigma \cap B_{\alpha^{-1}}$ with $\text{inj}_{\Sigma}(p_0) \leq \Delta \alpha^{-1}$.*

We can now prove our desired compactness result.

Proof. (Theorem 1.3) By Lemma 3.5, the injectivity radius of the sequence Σ_i is uniformly bounded below by $\delta_0 > 0$. Moreover, by Corollary 3.7 there is a point p_i in the ball $B_{\alpha^{-1}}(0)$ so that $\text{inj}_{\Sigma_i}(p_i) \leq \Delta \alpha^{-1}$. As a consequence, we may apply Theorem 1.4 and obtain a sub-sequence of the Σ_i that converges, with multiplicity one, uniformly in C^{∞} on compact subsets of $\mathbb{R}^3$ to some complete, embedded non-simply connected minimal surface Σ_{∞} . The inner and outer control on the genus and topology of the Σ_i carry over to the limit in the indicated manner. $\square$

4. APPLICATIONS

The compactness result developed in the previous section is particularly strong for sequences of genus-one surfaces, as there is only one scale for the genus. Indeed, there are several interesting scales for genus- g helicoids, the scale of the asymptotic helicoid and the scale(s) of the genus. In principle, one might expect a relationship between them. This is the case for genus one; however it is not clear there is such a connection for $g \geq 2$ – this is discussed in more detail in [1].

4.1. Compactness of $\mathcal{E}(1, 1)$. Let us now focus on the space $\mathcal{E}(1, 1)$, i.e. genus-one helicoids. To prove Theorem 1.1, we show that for any $\Sigma \in \mathcal{E}(1, 1)$, there is an upper and lower bound on the ratio between the scale of the genus and the scale of the asymptotic helicoid. To verify this bound and that the limit is also asymptotic to H , we must make a connection between the local nature of the convergence and global nature of the asymptotic scale. This is made by evaluating certain path integrals of holomorphic Weierstrass data. Indeed, we first establish a uniform R such that all vertical normals of each Σ_i (after a rotation) occur in $B_R(0)$. We then find annular ends Γ_i , conformally mapped to the same domain in $\mathbb{C}$ by $z_i = (x_3)_i + \sqrt{-1}(x_3^*)_i$ and with Weierstrass data as described in Corollary 1.2 of [2]. Finally, we use calculus of residues on $\partial\Gamma_i$ to establish uniform control on the center and radius of the genus for a sub-sequence.

Note that Theorem 1.1 of [2] implies that for $\Sigma \in \mathcal{E}(1, 1)$, there are two points where the Gauss map points parallel to the axis of the asymptotic helicoid. Thus, after translating, rescaling and rotating, any $\Sigma \in \mathcal{E}(1, 1)$ will satisfy the conditions of the following lemma.

Lemma 4.1. *Suppose $\Sigma \in \mathcal{E}(1, 1)$, the genus of Σ is centered at 0, $r(\Sigma) = 1$, and that g , the usual stereographic projection of the Gauss map of Σ , has only a single pole and single zero. Then there is an $R \geq 1$ independent of Σ so that the pole and zero of g lie in $B_R \cap \Sigma$.*

Proof. Suppose this was not the case. That is, one has a sequence $\Sigma_i \in \mathcal{E}(1, 1)$ so that the genus of each Σ_i is centered at 0, $r(\Sigma_i) = 1$, and each g_i has a single pole and single zero, at least one of which does not lie in B_i . By rotating Σ_i , we may assume the pole does not lie in B_i . By Theorem 1.3, a sub-sequence of the Σ_i converge uniformly in C^∞ on compact subsets of $\mathbb{R}^3$, with multiplicity 1, to $\Sigma_\infty \in \mathcal{E}(1, 1)$ where $r(\Sigma_\infty) = 1$.

Now consider $g_\infty : \Sigma_\infty \rightarrow \mathbb{C}$; it has at least one zero. Suppose g_∞ has more than one zero and call two such zeros $q_1, q_2 \in \Sigma_\infty$. For δ_0 as in Lemma 3.5, denote by σ_∞^j a closed, embedded curve in the component of $B_{\delta_0/2}(q_j) \cap \Sigma_\infty$ that contains q_j , chosen so that σ_∞^j surrounds q_j but neither surrounds nor contains any other pole or zero of g_∞ . Additionally, we choose the two σ_∞^j to be disjoint. Thus, $\int_{\sigma_\infty^j} \frac{dg_\infty}{g_\infty} = 2\pi\sqrt{-1}$.

By the convergence, there are simple, closed curves σ_i^j ($j = 1, 2$) in Σ_i so that σ_i^j converges smoothly to σ_∞^j . Let $D_i^j \subset \Sigma_i$ denote a disk such that $\partial D_i^j = \sigma_i^j$. For large enough i , $D_i^1 \cap D_i^2 = \emptyset$. There is at most one zero of g_i in $\cup_{j=1,2} D_i^j$ and the pole of g_i lies outside B_i . Thus, there exists i' such that, for $i \geq i'$, either $\int_{\sigma_i^1} \frac{dg_i}{g_i} = 0$ or the same is true for σ_i^2 . Letting $i \rightarrow \infty$ gives a contradiction; so g_∞ has only one zero. By Proposition 4.2 of [2], g_∞ has only one pole.

To conclude the proof, note there is an R_1 such that the pole, p_∞ , of g_∞ is in $B_{R_1} \cap \Sigma_\infty$. Let $\sigma_\infty \subset \Sigma_\infty$ be a smooth, embedded, closed curve in $B_{\delta_0/2}(p_\infty)$ that surrounds p_∞ and neither contains nor surrounds the zero of g_∞ . Thus, $\int_{\sigma_\infty} \frac{dg_\infty}{g_\infty} = -2\pi\sqrt{-1}$. Then by our convergence result, there are smooth, embedded, closed curves $\sigma_i \subset \Sigma_i \cap B_{3\delta_0/4}(p_\infty)$, with σ_i converging to σ_∞ . Note σ_i is necessarily null-homotopic and for i large enough so $\sigma_\infty \subset B_{i/2}(0)$ we compute $\frac{1}{2\pi\sqrt{-1}} \int_{\sigma_i} \frac{dg_i}{g_i} \geq 0$. As $i \rightarrow \infty$ the smooth convergence provides a contradiction. $\square$

Before proving, Theorem 1.1, we establish the necessary uniform control on the center and radius of the genus for surfaces in $\mathcal{E}(1, 1)$ that are asymptotic to a fixed helicoid H . Throughout the following proof, we make repeated use of [2].

Lemma 4.2. *Let $\Sigma_i \in \mathcal{E}(1, 1)$ and suppose that all the Σ_i are asymptotic to the same helicoid, H , which has axis the x_3 -axis. Then, there exist $C_1, C_2 > 0$ and a sub-sequence such that $1/C_1 \leq r(\Sigma_i) \leq C_1$ and, after a rotation, $|x_1(p_i)| + |x_2(p_i)| \leq C_2$, where p_i is the center of the genus of each Σ_i .*

Proof. Translate each Σ_i by $-p_i$ so that the genus of each of the Σ_i is centered at 0. Then rescale each Σ_i by α_i so that after the rescaling $r(\Sigma_i) = 1$. Thus, each rescaled and translated surface Σ_i is asymptotic to the helicoid $H_i = \alpha_i(H - p_i)$. By Theorem 1.3, passing to a sub-sequence, Σ_i converges uniformly on compact sets to $\Sigma_\infty \in \mathcal{E}(1, 1)$ with multiplicity one. By Corollary 1.3 of [2], Σ_∞ is asymptotic to some helicoid H_∞ . Since each Σ_i is asymptotic to H_i , which has vertical axis, the stereographic projection g_i of the Gauss map of Σ_i has exactly one pole and one zero. Thus, as in the proof of Lemma 4.1, we can conclude the same for Σ_∞ .

As each Σ_i satisfies the hypotheses of Lemma 4.1, there exists an R such that both the zero and pole of g_i ($i \leq \infty$) lie on the component, Σ_i^0 , of $B_R(0) \cap \Sigma_i$ containing the genus. Thus, $\Gamma_i = \Sigma_i \setminus \Sigma_i^0$ is an annulus and g_i has no poles or zeros in Γ_i . Hence, by the arguments of Section 4 of [2], $z_i = (x_3)_i + \sqrt{-1}(x_3)_i^* : \Gamma_i \rightarrow \mathbb{C}$ and $f_i = \log g_i : \Gamma_i \rightarrow \mathbb{C}$ are well-defined. (Here $(x_3)_i$ is x_3 restricted to Σ_i and $(x_3)_i^*$ is the harmonic conjugate of $(x_3)_i$.) Corollary 1.2 of [2] tells us that $f_i(p) =$

$\sqrt{-1}\lambda_i z_i(p) + F_i(p)$ where λ_i determines the scale of H_i and $F_i(p)$ is holomorphic on Γ_i and extends holomorphically to ∞ with $F(\infty) = 0$.

Let $A_C = \{z \in \mathbb{C} : |z| > C\}$. By suitably translating $(x_3)_i^*$, $A_{C/2} \subset z_\infty(\Gamma_\infty)$ for $C > 0$; thus there exists i_0 such that, for $i_0 \leq i \leq \infty$, $A_C \subset z_i(\Gamma_i)$. By precomposing with z_i^{-1} , we may think of f_i as a holomorphic function on A_C . If u is the standard coordinate of $\mathbb{C}$ restricted to A_C , then for $i_0 \leq i \leq \infty$, $f_i(u) = \sqrt{-1}\lambda_i u + F_i(u)$ where F_i is holomorphic and extends holomorphically and with a zero to ∞ . There is an $R' > R$ so that $\{p \in \Gamma_\infty : |z_\infty(p)| \leq 2C\} \subset B_{R'}$; thus, by perhaps increasing i_0 , for $i_0 \leq i \leq \infty$, $\{p \in \Gamma_i : |z_i(p)| \leq 2C\} \subset B_{2R'}$. As a consequence, the uniform convergence of Σ_i to Σ_∞ implies that for $\gamma = \{u : |u| = \frac{3}{2}C\} \subset A_C$, $f_i \rightarrow f_\infty$ in $C^\infty(\gamma)$. On the other hand, the calculus of residues implies that for $i_0 \leq i \leq \infty$,

$$(4.1) \quad \int_\gamma f_i(u) \frac{du}{u^2} = -2\pi\lambda_i$$

and hence we see immediately that $\lambda_i \rightarrow \lambda_\infty > 0$. Since the initial helicoid H had some λ_H associated with its Weierstrass data, this gives an upper and lower bound on the rescaling of each initial Σ_i , thus producing the necessary C_1 .

Since each F_i is holomorphic on A_C with a holomorphic extension to ∞ (and a zero there), we can expand in a Laurent series, i.e. for every $u \in A_C$ one has

$$(4.2) \quad F_i(u) = \sum_{j=1}^{\infty} \frac{a_{i,j}}{u^j}$$

where this is a convergent sum. Thus, for $i_0 \leq i \leq \infty$

$$(4.3) \quad \int_\gamma f_i(u) u^{j-1} du = 2\pi\sqrt{-1}a_{i,j}$$

and hence $\lim_{i \rightarrow \infty} a_{i,j} = a_{\infty,j}$. Thus, $F_i \rightarrow F_\infty$ uniformly in $\{u : 2C \leq |u| \leq \infty\}$.

The Weierstrass representation implies that H_i converge to H_∞ and hence that $x_1(p_i) \rightarrow x_1(p_\infty)$ and $x_2(p_i) \rightarrow x_2(p_\infty)$. This produces the necessary bound, C_2 , on the distance between the center of the genus and the axis of the initial helicoid H . Indeed, the convergence $F_i \rightarrow F_\infty$ implies there is a uniform C_0 so that for $u > 2C$ and $i_0 \leq i \leq \infty$, one can write $F_i(u) = \frac{a_{i,1}}{u} + \tilde{F}_i(u)$, where $|\tilde{F}_i(u)| \leq \frac{C_0}{u^2}$ and $|a_{i,1}| \leq C_0$. Recall the Weierstrass representation gives that

$$(4.4) \quad dx_1 - \sqrt{-1}dx_2 = g^{-1}dh - \bar{g}d\bar{h},$$

where $dh = dx_3 + \sqrt{-1}dx_3^*$ is the height differential. Integrating this form along $\{t + (\sqrt{-1})0 : t \in [3C, t_1]\} \subset A_C$, gives:

$$(4.5) \quad ((x_1)_i(t_1) - (x_1)_i(3C)) - \sqrt{-1}((x_2)_i(t_1) - (x_2)_i(3C)) = \int_{3C}^{t_1} J_i(t) dt$$

where $J_i(t) = e^{-\sqrt{-1}(\lambda_i t + \text{Im } F_i(t))}(e^{-\text{Re } F_i(t)} - e^{\text{Re } F_i(t)})$. For C sufficiently large, $|F_i(t)| \leq 1/2$ and so expanding in a power series,

$$(4.6) \quad J_i(t) = -2e^{-\sqrt{-1}\lambda_i t} \frac{\text{Re } a_{i,1}}{t} + G_i(t)$$

where here $|G_i(t)| \leq \frac{10C_0}{t^2}$.

The first term is a convergent (as $t_1 \rightarrow \infty$) oscillating integral while the integral of $G_i(t)$ is absolutely convergent. Thus, there exists a C_2 depending only on C_0

(and in particular independent of t_1, i) so, for $i_0 \leq i \leq \infty$, on each Σ_i ,

$$(4.7) \quad |(x_1)_i(t_1) - (x_1)_i(3C)| + |(x_2)_i(t_1) - (x_2)_i(3C)| \leq C_2.$$

Recall the translated surfaces Σ_i have genus centered at zero. Since one finds the axis by letting $t_1 \rightarrow \infty$, the above bound shows each translated surface has axis a uniform distance from the origin. Thus, the original center for each genus, p_i , satisfies the desired uniform bound. $\square$

The previous lemma tells us that for any sequence of surfaces in $\mathcal{E}(1, 1)$, asymptotic to a fixed helicoid, there is a sub-sequence with the scale of the genus uniformly controlled and the center of the genus lying inside of a cylinder. With this information, we can now show Theorem 1.1:

Proof. By the previous lemma, we know that there exists an $R > 0$ and a sub-sequence Σ_i such that $r(\Sigma_i) \rightarrow r_\infty > 0$ and the genus of these Σ_i is centered in a cylinder of radius R . Translate the surface parallel to the axis of the cylinder by some v_i so the genus is centered in $B_R(0)$. Use Theorem 1.3 to show that there is a sub-sequence so $\Sigma_i - v_i \rightarrow \Sigma_\infty \in \mathcal{E}(1, 1)$, which is asymptotic to some helicoid. The integral (4.1) implies this helicoid is H . If $\lim_{i \rightarrow \infty} |v_i| < \infty$ then $\Sigma_\infty \in \mathcal{E}(1, 1)$, and if $\lim_{i \rightarrow \infty} |v_i| = \infty$ then $\Sigma_\infty = H$. $\square$

4.2. Geometric Structure of $\mathcal{E}(1, 1, R)$. Theorem 1.1 of [3] gave an effective description of disks with large curvature by comparing them with helicoids. Theorem 1.3 allows us to do the same for elements $\mathcal{E}(1, 1, R)$.

Theorem 4.3. *Given $\epsilon > 0$ and $R \geq 1$ there exists an $R' = R'(\epsilon, R) \geq R$ so that: if $\Sigma \in \mathcal{E}(1, 1, R')$ with $r(\Sigma) = 1$ and the genus of Σ is centered at 0, then the component of $B_R(0) \cap \Sigma$ containing the genus is bi-Lipschitz with a subset of an element of $\mathcal{E}(1, 1)$ and the Lipschitz constant is in $(1 - \epsilon, 1 + \epsilon)$.*

Proof. The proof follows by contradiction exactly as in the proof of Theorem 1.1 of [3] with two very minor differences. First, Theorem 1.3 of this paper replaces the convergence to Σ_∞ outlined in that proof. Second, by Theorem 1.1 of [2], $C = \sup_{\Sigma_\infty} |A| < \infty$ and so for any ϵ , we find a smooth ν_i defined on a subset of Σ_∞ so that $C|\nu_i| + |\nabla_{\Sigma_\infty} \nu_i| < \epsilon$ and the graph of ν_i is the component of $\Sigma_i \cap B_{R'}$ containing the genus. For details, see Section 6 of [3]. $\square$

APPENDIX A. THE LAMINATION THEORY OF [5]

In [5], Colding and Minicozzi generalize their lamination theory for minimal disks – i.e. Theorem 0.1 of [9] – to arbitrary sequences of minimal surfaces of fixed, finite genus. To do so, they must allow for a more general class of singular models, as is clear from considering a rescaling of the catenoid. Surfaces modeled on the neck of a catenoid form an important class, one that is characterized by having genus zero and disconnected boundary. Another important class, especially for our purposes, are surfaces of finite genus and connected boundary. This second class of surfaces, because they have connected boundary, turn out to be structurally very similar to disks. Indeed, most of the results of [6, 7, 8, 9] hold for them (in a suitably form) and with only slight modifications of the proofs; this has important ramifications for lamination theory.

We first briefly discuss the most general lamination result of [5] – Theorem 0.14. Suppose Σ_i is a sequence of embedded minimal surfaces with fixed, finite genus and

$\partial\Sigma_i \subset B_{R_i}$ with $R_i \rightarrow \infty$. Colding and Minicozzi show that if the curvature of the sequence blows up at a point $y \in \mathbb{R}^3$ (i.e. if for all $r > 0$, $\sup_i \sup_{B_r(y) \cap \Sigma_i} |A|^2 = \infty$), then after a rotation, a sub-sequence Σ_i converge to the singular lamination $\mathcal{L} \setminus \mathcal{S}$ in the C^α topology ($\alpha \in (0, 1)$) and the curvature blows precisely at points of $\mathcal{S}$. Here $\mathcal{L} = \{x_3 = t\}_{t \in \mathcal{I}}$, $\{x_3(y) : y \in \mathcal{S}\} = \mathcal{I}$ a closed subset of $\mathbb{R}^3$. Notice little global information about the limit lamination is given; this contrasts with the case for disks where $\mathcal{I} = \mathbb{R}$ and $\mathcal{S}$ is a Lipschitz graph over the x_3 -axis – in fact a line.

More generally, the topology of the sequence can restrict $\mathcal{I}$ and give more information about convergence near $\mathcal{S}$ and the structure of $\mathcal{S}$. To make this precise, we distinguish between two types of singular points $y \in \mathcal{S}$. Heuristically, the distinction is between points where on small scales near the point all the Σ_i are disks and points where on small scales near the point all the Σ_i contain necks. This is the exact description if the genus of the surfaces is zero, but must be refined for sequences with positive genus. Following [5], we make this precise for a sequence Σ_i converging to a lamination $\mathcal{L}$ with singular set $\mathcal{S}$:

Definition A.1. We say $y \in \mathcal{S}$ is an element of $\mathcal{S}_{ulsc}$ if there exist both $r > 0$ fixed and a sequence $r_i \rightarrow 0$ such that $B_r(y) \cap \Sigma_i$ and $B_{r_i}(y) \cap \Sigma_i$ have the same genus and every component of $B_{r_i}(y) \cap \Sigma_i$ has connected boundary.

Definition A.2. We say $y \in \mathcal{S}$ is an element of $\mathcal{S}_{neck}$ if there exist both $r > 0$ fixed and a sequence $r_i \rightarrow 0$ such that $B_r(y) \cap \Sigma_i$ and $B_{r_i}(y) \cap \Sigma_i$ have the same genus and $B_{r_i}(y) \cap \Sigma_i$ has at least one component with disconnected boundary.

For instance, if the Σ_i are the homothetic blow-down of a helicoid or genus-one helicoid, then 0 is an element of $\mathcal{S}_{ulsc}$, whereas if the Σ_i are the homothetic blow-down of a catenoid then 0 is an element of $\mathcal{S}_{neck}$. Colding and Minicozzi show that near a point of $\mathcal{S}_{ulsc}$ or $\mathcal{S}_{neck}$ these examples model the behavior of the convergence.

A major result of [5] refines the general compactness theorem based on more careful analysis of the topology of the sequence. One important example is the no-mixing theorem (i.e. Theorem 0.4 of [5]), which states that, up to a passing to a sub-sequence, either $\mathcal{S} = \mathcal{S}_{ulsc}$ or $\mathcal{S} = \mathcal{S}_{neck}$. This is particularly important as ULSC sequences (i.e. sequences where $\mathcal{S} = \mathcal{S}_{ulsc}$) behave much like disks. Indeed, Theorem 0.9 of [5] states:

Theorem A.3. *Let $\Sigma_i \subset B_{R_i} = B_{R_i}(0) \subset \mathbb{R}^3$ be a sequence of compact embedded minimal surfaces of fixed genus with $\partial\Sigma_i \subset \partial B_{R_i}$ where $R_i \rightarrow \infty$. Assume the sequence Σ_i is ULSC. If $\sup_{B_1 \cap \Sigma_i} |A|^2 \rightarrow \infty$, then there exists a sub-sequence Σ_j , a foliation $\mathcal{L} = \{x_3 = t\}_t$ of $\mathbb{R}^3$ by parallel planes, and a closed nonempty set $\mathcal{S}$ in the union of the leaves of $\mathcal{L}$ such that after a rotation of $\mathbb{R}^3$:*

- (1) *For each $\alpha \in (0, 1)$, $\Sigma_j \setminus \mathcal{S}$ converges in the C^α -topology to $\mathcal{L} \setminus \mathcal{S}$.*
- (2) *$\sup_{B_r(x) \cap \Sigma_j} |A|^2 \rightarrow \infty$ as $j \rightarrow \infty$ for all $r > 0$ and $x \in \mathcal{S}$.*
- (3) *$\mathcal{S}$ is orthogonal to the leaves of the lamination and is either the union of two disjoint lines $\mathcal{S}_1$ and $\mathcal{S}_2$, or is a single line.*
- (4) *Away from $\mathcal{S}$, each Σ_j consists of exactly two multi-valued graphs spiraling together. Near $\mathcal{S}$ the pair of multi-valued graphs form double spiral staircases with opposite orientations. Thus, circling only one component of $\mathcal{S}$ results in going either up or down. If $\mathcal{S} = \mathcal{S}_1 \cup \mathcal{S}_2$ then the multi-valued graphs are glued so that a path circling both $\mathcal{S}_1$ and $\mathcal{S}_2$ closes up.*

Remark A.4. The situation with two singular sets is modeled on the degeneration of the Riemann examples. As such the limit can be thought of as having “infinite” topology. This is made rigorous by noting that for $R > 0$ large enough so $B_{R/2}(0)$ intersects both singular lines, then (4) implies that for any $f \in \mathbb{Z}^+$ for i sufficiently large $B_R(0) \cap \Sigma_i$ has a component with at least f boundary components.

REFERENCES

1. J. Bernstein and C. Breiner, *Compactness Properties of the space of genus- g helicoids*, Preprint.
2. ———, *Conformal structure of minimal surfaces with finite topology*, To Appear, Comm. Math. Helv. <http://arxiv.org/abs/0810.4478>.
3. ———, *Helicoid-like minimal disks and uniqueness*, Preprint. <http://arxiv.org/abs/0802.1497>.
4. H. I. Choi and R. Schoen, *The space of minimal embeddings of a surface into a three-dimensional manifold of positive Ricci curvature*, Inventiones Mathematicae **81** (1985), no. 3, 387–394.
5. T. H. Colding and W. P. Minicozzi II, *The Space of Embedded Minimal Surfaces of Fixed Genus in a 3-manifold V; Fixed Genus*, Preprint. <http://arxiv.org/abs/math/0509647>.
6. ———, *The space of embedded minimal surfaces of fixed genus in a 3-manifold I; Estimates off the axis for disks*, Ann. of Math. (2) **160** (2004), no. 1, 27–68.
7. ———, *The space of embedded minimal surfaces of fixed genus in a 3-manifold II; Multi-valued graphs in disks*, Ann. of Math. (2) **160** (2004), no. 1, 69–92.
8. ———, *The space of embedded minimal surfaces of fixed genus in a 3-manifold III; Planar domains*, Ann. of Math. (2) **160** (2004), no. 2, 523–572.
9. ———, *The space of embedded minimal surfaces of fixed genus in a 3-manifold IV; Locally simply connected*, Ann. of Math. (2) **160** (2004), no. 2, 573–615.
10. ———, *The Calabi-Yau conjectures for embedded surfaces*, Ann. of Math. **167** (2008), no. 1, 211–243.
11. D. Fischer-Colbrie and R. Schoen, *The structure of complete stable minimal surfaces in 3-manifolds of nonnegative scalar curvature*, Comm. Pure Appl. Math. **33** (1980), 199–211.
12. D. Hoffman and B. White, *Genus-one helicoids from a variational point of view*, Comm. Math. Helv. **83** (2008), no. 4, 767–813.
13. W. H. Meeks III, J. Perez, and A. Ros, *Embedded minimal surfaces: removable singularities, local pictures and parking garage structures, the dynamics of dilation invariant collections and the characterization of examples of quadratic curvature decay*, Preprint.
14. ———, *The Geometry of Minimal Surfaces of Finite Genus III; Bounds on the Topology and Index of Classical Minimal Surfaces*, Preprint.
15. W. H. Meeks III and H. Rosenberg, *The minimal lamination closure theorem*, Duke Math. J. **133** (2006), no. 3, 467–497.
16. R. Schoen, *Seminar on minimal submanifolds*, Ann. of Math. Studies, vol. 103, ch. Estimates for stable minimal surfaces in three-dimensional manifolds, pp. 111–126, Princeton University Press, Princeton, N.J., 1983.

DEPT. OF MATH, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02139, USA
E-mail address: jbern@math.mit.edu

DEPT. OF MATH, JOHNS HOPKINS UNIVERSITY, BALTIMORE, MD 21218, USA
E-mail address: cbreiner@math.jhu.edu